

Bodyweight (kg) of dog	LIBRELA strength (mg) to be administered				
	5mg	10mg	15mg	20mg	30mg
5.0 - 10.0	1 vial				
10.1 - 20.0		1 vial			
20.1 - 30.0			1 vial		
30.1 - 40.0				1 vial	
40.1 - 60.0					1 vial
60.1 - 80.0				2 vials	
80.1 - 100.0				1 vial	+ 1 vial
100.1 - 120.0					2 vials



Store in a refrigerator, between 2° to 8°C. Do not freeze.



Once opened, contents of the vial should be used immediately and any remaining solution should be disposed.

Each vial delivers 1 mL volume for injection.

For dogs weighing <5.0 kg: aseptically withdraw 0.1 mL/kg from a single 5 mg/mL vial and administer subcutaneously. For volumes ≤ 0.5 mL, use an adequate syringe size and dose to the nearest 0.1 mL. For dogs above 60 kg, the contents of more than one vial are required to administer a single dose. In those cases, withdraw the content from each required vial into the same syringe and administer as a single subcutaneous injection (2 mL).

Librela[®] contains Bedinvetmab. Librela is licensed for the alleviation of Pain associated with osteoarthritis in dogs. POM-V

For further information please see the product's SPC or contact Zoetis Singapore Pte Ltd, 108 Pasir Panjang Rd, #04-04/07 Golden Agri Plaza, Singapore 118535.
www.zoetis.sg - Customer Support: zoetissingapore@zoetis.com